F_B from moving B mesons

S.Collins[a], G.Cowan[a], C. T. H. Davies[a], J. Hein[b], G. P. Lepage[c], C. Morningstar[d] and J. Shigemitsu[e].

[a]Department of Physics and Astronomy, University of Glasgow, Glasgow, G12 8QQ, UK

[b]Department of Physics and Astronomy, University of Edinburgh, Edinburgh, EH9 3JZ, UK

[c]Newman Laboratory of Nuclear Science, Cornell University, Ithaca NY 14853, USA

[d]Department of Physics, Carnegie Mellon University, Pittsburgh PA 15213, USA

[e]Department of Physics, the Ohio State University, OH 43210 USA

arXiv:hep-lat/0110003v1 1 Oct 2001

We show results for the B meson decay constant calculated both for B mesons at rest and those with non-zero momentum and using both the temporal and spatial components of the axial vector current. It is an important check of lattice systematic errors that all these determinations of f_B should agree. We also describe how well different smearings for the B meson work at non-zero momentum - the optimal smearing has a narrow smearing on the b quark.

1. Introduction

Matrix elements involving moving B and D mesons are important for studies of $B \to D$ and $B \to \pi$ decay. It is necessary to understand the systematic errors in lattice QCD that come from the presence of non-zero momenta and the optimal way in which to handle such mesons on the lattice. An easy place to study these effects is in the determination of the B meson decay constant, f_B [1].

In the absence of discretisation errors it should be true that

$$< 0|A_\mu|B >= f_B p_\mu. \quad (1)$$

We use both A_0 and A_k with all current corrections and renormalisation through α_s/M_b [2].

Results shown are for an ensemble of 278 $12^3 \times 24$ lattices at $\beta = 5.7$. Clover light quarks are used with $\kappa = \kappa_s$ and NRQCD heavy quarks are used with masses in lattice units of 2 and 8 ($m_b a = 4.0$). All momenta are considered up to $(pa)^2$ of 16 in units of $(2\pi/L)^2$, i.e. roughly 4 GeV^2 at this lattice spacing.

2. Smearing

We generated heavy and light quark propagators with 3 different smearings: a delta function, a narrow Gaussian (width 1) and a broad Gaussian (width 3). We combined these smearings together to make 6 different smearings for the heavy-light meson and analysed the resulting smeared-smeared meson correlators using the constrained curve fitting methods described in [3]. The results are shown in Figure 1, which plots the amplitude of the ground state B meson as a function of squared momentum for some of these smearings. It is clear that the optimal smearing for a moving heavy-light meson is one in which the smearing on the heavy quark is a narrow one and the smearing on the light quark is a broad one. This is not surprising if one considers that the heavy quark is carrying almost all of the meson momentum. The overlap as a function of momentum is maximized if the heavy quark has a delta function smearing, but in that case the statistical noise is large because of the well-known problem that an unsmeared heavy quark has a poor signal/noise ratio [4]. At zero momentum this has led to the received wisdom that the heavy quark should have a broad smearing - Figure 1



Figure 1. The amplitude for the ground state in various heavy-light smeared-smeared correlators as a function of $\vec{p}^2$ in units of $(2\pi/L)^2$. The heavy mass was $ma = 8$.



Figure 2. The effective energy splitting between smeared-smeared correlators of $\vec{p}^2 = 0$ and 2 in units of $(2\pi/L)^2$ against time, for smearings 3 and 5 of Figure 1.

shows that this is not correct at non-zero momentum.

Figure 2 amplifies this point by showing the rapid plateau and consequent small error on the kinetic energy possible with a good finite-momentum smearing, in contrast to that obtained with a smearing that might have been considered a good one at zero momentum.

3. f_B from A_0

We used smearing '5' from the above analysis and the constrained curve fitting methods to determine f_B for moving B mesons and the temporal axial current. Matrix elements for A_0 are obtained from a simultaneous fit to *all* A_0^i (i=0,1,2) and A_k^i (i=0,1,2,3,4) current correlators at a given momentum. The construction of the continuum A_0 from $A_0^{i,latt}$ is described in [2].

Figure 3 shows results for the ratio:

$$R(p) = \frac{\langle 0|A_0|B(p)\rangle/\sqrt{E}}{\langle 0|A_0|B(0)\rangle/\sqrt{M}} \qquad (2)$$



Figure 3. The ratio $R(p)$ (see text) for A_0 as a function of momentum, pa. The curve is $\sqrt{E/M}$ where E is the heavy-light meson energy and M its mass.



Figure 4. The ratio $R_k(p)$ (see text) for A_k as a function of momentum, $p_k a$.

for ma=8 using $\alpha_s(2/a)$ in the renormalisation. The expected curve $\sqrt{E/M}$ is also shown. No disagreement is seen until the highest momentum. At $ma = 2$, larger discrepancies appear [5].

4. f_B from A_k

The matrix elements for the spatial axial current behave rather differently to those for the temporal axial current since they must vanish when there is no component of $\vec{p}$ along the direction of the current. Even the leading order current, $A_k^0 = \overline{q}\gamma_5\gamma_k Q$ then has a matrix element $\mathcal{O}(p_k/M)$ with respect to A_0^0. This is of the same order as the '$1/M$' suppressed current contributions from $A_k^1 = \overline{q}\gamma_5\gamma_k(\vec{\gamma}\cdot\vec{D}/2M)Q$ and $A_k^3 = \overline{q}\gamma_5(\vec{D}/2M)Q$. A_k^1 contributes at tree level and A_k^3 at one-loop to the final result for A_k. To understand the size of different current contributions it is important to use a power-counting in both the external velocity and in Λ_{QCD}/M [5].

Figure 4 shows results for the ratio:

$$R_k(p) = \frac{\langle 0|A_k|B(p)\rangle}{\langle 0|A_0|B(p)\rangle} \qquad (3)$$

as a function of p_k for a range of values of $\vec{p}^2$, for ma=8. The line shown is p_k/E (variation with $\vec{p}^2$ of this line is not significant). Good agreement with the line is found for $p_k a$ = 1,2 ($\times 2\pi/L$) at all $\vec{p}^2$ but $p_k a$ = 3,4 show signs of deviation, presumably from missing current corrections that are higher order in the external velocity.

Figure 4 shows agreement between f_B from A_0 and A_k which resolves a long-standing problem. Ref. [1] found a disagreement of $\mathcal{O}(10\%)$ when comparing the tree-level matrix elements for clover fermions. This ignores the contribution of A_k^3 and does not allow for the clover equivalent of the different renormalisation of $A_{0,k}^0$ and $A_{0,k}^1$. Both these effects are included here.

5. Conclusions

We have demonstrated the usefulness of optimal (i.e. narrow) smearing for heavy quarks in moving heavy-light mesons, and of constrained curve fitting in extracting results out to much higher momenta than have previously been attempted. Good agreement is found for f_B at zero and non-zero momentum. For the temporal axial current at large ma this holds out to the highest momenta studied. To understand the behaviour of the currents at non-zero momentum it is important to use a power-counting appropriate to moving heavy-light mesons.

Acknowledgements SC, GC, CD and JH are members of the UKQCD collaboration. We are grateful to the following bodies for support of this work: PPARC, DoE, NSF and the EU under HPRN-2000-00145 Hadrons/Lattice QCD.

REFERENCES

1. J. Simone, Nucl. Phys. B (Proc. Suppl. **53**) 1997 386.
2. C. Morningstar and J. Shigemitsu, PRD**59** (1999) 094504.
3. G. P. Lepage, these Proceedings.
4. G. P. Lepage, NPB (Proc. Suppl. **26**) 45 1992.
5. S. Collins *et al*, in preparation.